

Excitatory neuronal role of PLCG2 in Alzheimer's disease

Tae Yeon Kim¹, Diana Acosta¹, Nicholas Sweeney¹, Marie-Amandine Bonte¹, Qi Guo², Ethan Rose¹, Jean Paul Chadarevian³, Hiroyasu Ogawa⁴, Tomohiro Kurosaki⁵, Qin Ma², Hayk Davtyan³, Mathew Blurton-Jones³, and **Hongjun Fu**^{1,6,*}

¹ Department of Neuroscience, The Ohio State University, Columbus, Ohio, USA

² Department of Biomedical Informatics, The Ohio State University, Columbus, Ohio, USA

³ UCI Institute for Memory Impairments and Neurological Disorders, University of California, Irvine, California, USA

⁴ Department of Orthopaedic Surgery, Gifu University Graduate School of Medicine, Gifu, Japan

⁵ Laboratory of Lymphocyte Differentiation, WPI Immunology Frontier Research Center, Osaka University, Osaka, Japan

⁶ Chronic Brain Injury Program, The Ohio State University, Columbus, Ohio, USA

* Corresponding Authors: H.F. (Hongjun.Fu@osumc.edu)

Abstract

Recent Genome-Wide Association Studies (GWAS) have identified novel rare coding variants (P522R, M28L variants) of the enzyme phospholipase-C- γ 2 (PLCG2) in late-onset Alzheimer's Disease (LOAD). The role of PLCG2 and its variants in AD has been largely investigated in microglia and responses to A β pathology; however, little is known about the role of PLCG2 and its AD variants in other cell types or its role in tau pathology. To evaluate the immunoreactivity of PLCG2 and its relationship with tau pathology, we performed western blot and immunofluorescence staining on human AD post-mortem brains and PS19 tau mice. By stereotaxic injection of AAV8-CaMKIIa-cre and AD TBS-soluble tau seeds in PLCG2-floxed mice, we quantified tau spreading induced by PLCG2 knockdown. Furthermore, we generated human neural organoids derived from human iPSCs of a healthy donor and a CRISPR/Cas9-edited PLCG2 P522R knock-in (KI) cell line to investigate the autophagy-lysosomal pathway (ALP) dynamics induced by PLCG2 by using FUW-mCherry-GFP-LC3 reporter assay. We found higher global protein levels of PLCG2 in human AD cases and PS19 mice, but pathological tau-positive cells had a lower level of PLCG2 than neighboring pathological tau-negative cells. Knockdown of PLCG2 in excitatory neurons increased tau spreading in PLCG2-floxed mice. Autophagy flux increased in P522R KI organoids, even in the presence of Bafilomycin A1 (autophagy flux inhibitor), compared to wild-type organoids. Our results suggest that repression or dysfunction of PLCG2 may contribute to tau pathology in excitatory neurons of AD, probably via the regulation of the ALP.

Sponsored By: BrightFocus Alzheimer's Disease Research Award, A2025025S

Presenter Name and contact information:

Hongjun "Harry" Fu, Ph.D., Associate Professor
Department of Neuroscience
The Ohio State University Wexner Medical Center
Columbus, Ohio, USA
Email: Hongjun.Fu@osumc.edu